arXiv:1608.03579v1 [math.RT] 11 Aug 2016

FINITE GENERATION OF COHOMOLOGY OF FINITE GROUPS

RAPHAËL ROUQUIER

Abstract. We give a proof of the finite generation of the cohomology ring of a finite p-group over $\mathbf{F}_p$ by reduction to the case of elementary abelian groups, based on Serre's Theorem on products of Bocksteins.

1. Definitions and basic properties

Let $k = \mathbf{F}_p$. Given G a finite group, we put $H^*(G) = H^*(G, k)$. We refer to [Ev] for results on group cohomology.

Given A a ring and M an A-module, we say that M is finite over A if it is a finitely generated A-module.

Let G be a finite group and L a subgroup of G. We have a restriction map $\mathrm{res}^G_L : H^*(G) \to H^*(L)$. It gives $H^*(L)$ the structure of an $H^*(G)$-module.

We denote by $\mathrm{norm}^G_L : H^*(L) \to H^*(G)$ the norm map. If L is central in G, then we have $\mathrm{res}^G_L \mathrm{norm}^G_L(\xi) = \xi^{[G:L]}$ for all $\xi \in H^*(L)$.

When L is normal in G, we denote by $\mathrm{inf}^G_{G/L} : H^*(G/L) \to H^*(G)$ the inflation map.

Let E be an elementary abelian p-group. The Bockstein $H^1(E) \to H^2(E)$ induces an injective morphism of algebras $S(H^1(E)) \hookrightarrow H^*(E)$. We denote by $H^*_{\mathrm{pol}}(E)$ its image. Note that $H^*(E)$ is a finitely generated $H^*_{\mathrm{pol}}(E)$-module and given $\xi \in H^*(E)$, we have $\xi^p \in H^*_{\mathrm{pol}}(E)$.

2. Finite generation for finite groups

The following result is classical. We provide here a proof independent of the finite generation of cohomology rings.

Lemma 2.1. *Let G be a p-group and E an elementary abelian subgroup. Then, $H^*(E)$ is finite over $H^{\mathrm{even}}(G)$.*

Proof. The result is straightforward when G is elementary abelian. As a consequence, given G, it is enough to prove the lemma when E is a maximal elementary abelian subgroup. We prove the lemma by induction on $|G|$. Let $Z \leq Z(G)$ with $|Z| = p$. Let P be a complement to Z in E. Let $A = \mathrm{inf}^E_{E/Z}(H^*_{\mathrm{pol}}(E/Z))$. Let x be a generator of $H^2(Z) \xrightarrow{\sim} H^2(E/P)$ and $y = \mathrm{inf}^E_{E/P}(x)$. We have

$$H^*_{\mathrm{pol}}(E) = A \otimes k[y].$$

Let $\xi = \mathrm{res}^G_E(\mathrm{norm}^G_Z(x)^p)$. We have $\mathrm{res}^E_Z(\xi) = x^{p[G:Z]}$, so $\xi - y^{p[G:Z]} \in H^*_{\mathrm{pol}}(E) \cap \ker \mathrm{res}^E_Z = A^{>0} H^*_{\mathrm{pol}}(E)$. We deduce that $H^*(E)$ is finite over its subalgebra generated by A and ξ.

Date: October 2014.

The author was partially supported by the NSF grant DMS-1161999.

By induction, $H^*(E/Z)$ is finite over $H^*(G/Z)$. We deduce that $H^*(E)$ is finite over its subalgebra generated by ξ and $\inf_{E/Z}^E \operatorname{res}_{E/Z}^{G/Z} H^*(G/Z) = \operatorname{res}_E^G \inf_{G/Z}^G H^*(G/Z)$. □

Let us recall a form of Serre's Theorem on product of Bocksteins [Se]. We state the result over the integers for a useful consequence stated in Corollary 2.3.

Theorem 2.2 (Serre)**.** *Let G be a finite p-group. Assume G is not elementary abelian. Then, there is $n \geq 2$, there are subgroups $H_1, \dots, H_n$ of index p of G and an exact sequence of $\mathbf{Z}G$-modules*

$$0 \to \mathbf{Z} \to \operatorname{Ind}_{H_n}^G \mathbf{Z} \to \cdots \to \operatorname{Ind}_{H_1}^G \mathbf{Z} \to \mathbf{Z} \to 0$$

defining a zero class in $\operatorname{Ext}^n_{\mathbf{Z}G}(\mathbf{Z}, \mathbf{Z})$.

Proof. Serre shows there are elements $z_1, \dots, z_m \in H^1(G, \mathbf{Z}/p)$ such that $\beta(z_1) \cdots \beta(z_m) = 0$. The element z_i corresponds to a surjective morphism $G \to \mathbf{Z}/p$ with kernel H_i, and we identify $\operatorname{Ind}_{H_i}^G \mathbf{Z}$ with $\mathbf{Z}[G/H_i] = \mathbf{Z}[\sigma]/(\sigma^p - 1)$, where σ is a generator of G/H_i. The element $\beta(z_i) \in H^2(G, \mathbf{Z}/p)$ is the image of the class $c_i \in H^2(G, \mathbf{Z})$ given by the exact sequence

$$0 \to \mathbf{Z} \xrightarrow{1+\sigma+\cdots+\sigma^{p-1}} \operatorname{Ind}_{H_i}^G \mathbf{Z} \xrightarrow{1-\sigma} \operatorname{Ind}_{H_i}^G \mathbf{Z} \xrightarrow{\text{augmentation}} \mathbf{Z} \to 0.$$

Let $c = c_1 \cdots c_m \in H^{2m}(G, \mathbf{Z})$. The image of c in $H^{2m}(G, \mathbf{Z}/p)$ vanishes, hence $c \in pH^{2m}(G, \mathbf{Z})$. Fix r such that $|G| = p^r$. Since $|G| H^{>0}(G, \mathbf{Z}) = 0$, we deduce that $c^r = 0$. □

We will only need the case $R = \mathbf{F}_p$ of the corollary below. We denote by $D^b(RG)$ the derived category of bounded complexes of finitely generated RG-modules.

Corollary 2.3. *Let G be a finite group and R a discrete valuation ring with residue field of characteristic p or a field of charactetistic p. Assume $x^{p-1} = 1$ has $p-1$ solutions in R.*

Let $\mathcal{I}$ be the thick subcategory of $D^b(RG)$ generated by modules of the form $\operatorname{Ind}_E^G M$, where E runs over elementary abelian subgroups of G and M runs over one-dimensional representations of E over R.

We have $\mathcal{I} = D^b(RG)$.

Proof. Assume first G is an elementary abelian p-group. Let L be a finitely generated RG-module. Consider a projective cover $f : P \to L$ and let $L' = \ker f$. The R-module L' is free, so L' is an extension of RG-modules that are free of rank 1 as R-modules. So $L' \in \mathcal{I}$ and similarly $P \in \mathcal{I}$, hence $L \in \mathcal{I}$. As a consequence, the corollary holds for G elementary abelian.

Assume now G is a p-group that is not elementary abelian. We proceed by induction on $|G|$. Let L be a finitely generated RG-module. By induction, $\operatorname{Ind}_H^G \operatorname{Res}_H^G(L) \in \mathcal{I}$ whenever H is a proper subgroup of G. Applying $L \otimes_{\mathbf{Z}G} -$ to the exact sequence of Theorem 2.2, we obtain an exact sequence

$$0 \to L \to \operatorname{Ind}_{H_n}^G \operatorname{Res}_{H_n}^G(L) \to \cdots \to \operatorname{Ind}_{H_1}^G \operatorname{Res}_{H_1}^G(L) \to L \to 0.$$

Since that sequence defines the zero class in $\operatorname{Ext}^n(L, L)$, it follows that L is a direct summand of $0 \to \operatorname{Ind}_{H_n}^G \operatorname{Res}_{H_n}^G(L) \to \cdots \to \operatorname{Ind}_{H_1}^G \operatorname{Res}_{H_1}^G(L) \to 0$ in $D^b(RG)$. We deduce that $L \in \mathcal{I}$.

Finally, assume G is a finite group. Let P be a Sylow p-subgroup of G and let L a finitely generated RG-module. We know that $\operatorname{Ind}_P^G \operatorname{Res}_P^G(L) \in \mathcal{I}$. Since L is a direct summand of $\operatorname{Ind}_P^G \operatorname{Res}_P^G(L)$, we deduce that $L \in \mathcal{I}$. □

Remark 2.4. Corollary 2.3 implies a corresponding generation result for the stable category of RG. That was observed by the author in the mid 90s and communicated to J. Carlson who wrote an account of this in [Ca].

Theorem 2.5 (Golod, Venkov, Evens)**.** *Let G be a finite p-group. The ring $H^*(G)$ is finitely generated. Given M a finitely generated kG-module, then $H^*(G, M)$ is a finitely generated $H^*(G)$-module.*

Note that the case where G is an arbitrary finite group follows easily, cf [Ev].

Proof. Let S be a finitely generated subalgebra of $H^{\text{even}}(G)$ such that $H^*(E)$ is a finitely generated S-module for every elementary abelian subgroup E of G. Such an algebra exists by Lemma 2.1.

Let $\mathcal{J}$ be the full subcategory of $D^b(kG)$ of complexes C such that the S-module $H^*(G, C) = \bigoplus_i \mathrm{Hom}_{D^b(kG)}(k, C[i])$ is finitely generated.

Let $C_1 \to C_2 \to C_3 \rightsquigarrow$ be a distinguished triangle in $D^b(kG)$. We have a long exact sequence

$$\cdots \to H^i(C_1) \to H^i(C_2) \to H^i(C_3) \to H^{i+1}(C_1) \to \cdots$$

Assume $C_1, C_3 \in \mathcal{J}$. Let I be a finite generating set of $H^*(C_1)$ as an S-module and J a finite generating set of $\ker(H^*(C_3) \to H^{*+1}(C_1))$ as an S-module. Let I' be the image of I in $H^*(C_2)$ and let J' be a finite subset of $H^*(C_2)$ with image J. Then, $I' \cup J'$ generates $H^*(C_2)$ as an S-module, hence $C_2 \in \mathcal{J}$.

Note that if $C \oplus C' \in \mathcal{J}$, then $C \in \mathcal{J}$. We deduce that $\mathcal{J}$ is a thick subcategory of $D^b(kG)$.

Let E be an elementary abelian subgroup of G. Since $H^*(G, \mathrm{Ind}_E^G(k)) \simeq H^*(E, k)$ is a finitely generated S-module, we deduce that $\mathrm{Ind}_E^G(k) \in \mathcal{J}$.

We deduce from Corollary 2.3 that $\mathcal{J} = D^b(kG)$. □

References

[Ca] J. Carlson, *Cohomology and induction from elementary abelian subgroups*, Q. J. Math. **51** (2000), 169–181.
[Ev] L. Evens, "The cohomology of groups", Oxford University Press, 1991.
[Se] J.P. Serre, *Sur la dimension cohomologique des groupes profinis*, Topology **3** (1965), 413-420.

Department of Mathematics, UCLA, Box 951555, Los Angeles, CA 90095-1555, USA
E-mail address: rouquier@math.ucla.edu